



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C12N 15/82, A01H 5/00, 5/10 // C12N 15/53	A1	(11) International Publication Number: WO 99/53081 (43) International Publication Date: 21 October 1999 (21.10.99)
<p>(21) International Application Number: PCT/GB99/01059</p> <p>(22) International Filing Date: 7 April 1999 (07.04.99)</p> <p>(30) Priority Data: 9807818.1 9 April 1998 (09.04.98) GB</p> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): SHIPTON, Catherine, Ann [GB/GB]; Jealotts Hill Research Station, Bracknell, Berkshire RG42 6ET (GB). BRYAN, Ian, Bennett [GB/GB]; Jealotts Hill Research Station, Bracknell, Berkshire RG42 6ET (GB).</p> <p>(74) Agents: GAAL, Jozsef, Christopher et al.; Zeneca Agrochemicals, Intellectual Property Dept., P.O. Box 3538, Jealott's Hill Research Station, Bracknell, Berkshire RG42 6YA (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> </p>
<p>(54) Title: METHOD OF PRODUCING PLANTS WHICH ARE TOLERANT OR RESISTANT TO HERBICIDES</p>		
<p>(57) Abstract</p> <p>A method of making plants which are resistant or tolerant to herbicides which, <i>in vitro</i>, inhibit 4-hydroxyphenylpyruvate dioxygenase (4HPPD) comprises the steps of: (i) transforming plant material with a polynucleotide comprising a region encoding a phytoene desaturase; (ii) regenerating the thus transformed material into morphologically normal plants. In a preferred embodiment the region comprised by the polynucleotide is the sequence depicted in SEQ ID No.1, or is a sequence which is complementary to one which when incubated at a temperature of between 55 and 60 °C in 0.3 strength citrate buffered saline containing 0.1 % SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered saline containing 0.1 % SDS still hybridises with the sequence depicted in SEQ ID No.1.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CN	China	KR	Republic of Korea	PL	Poland		
CU	Cuba	KZ	Kazakhstan	PT	Portugal		
CZ	Czech Republic	LC	Saint Lucia	RO	Romania		
DE	Germany	LI	Liechtenstein	RU	Russian Federation		
DK	Denmark	LK	Sri Lanka	SD	Sudan		
EE	Estonia	LR	Liberia	SE	Sweden		
				SG	Singapore		

METHOD OF PRODUCING PLANTS WHICH ARE TOLERANT OR RESISTANT TO
HERBICIDES

The present invention relates *inter alia*, to a method of producing plants which are
5 tolerant or resistant to herbicides and in particular to the production of transgenic plants
which exhibit substantial resistance or substantial tolerance to herbicides when compared
with non transgenic like plants.

Plants which are substantially "tolerant" to a herbicide when they are subjected to it provide
a dose/response curve which is shifted to the right when compared with that provided by
10 similarly subjected non tolerant like plants. Such dose/response curves have "dose" plotted
on the x-axis and "percentage kill", "herbicidal effect" etc. plotted on the y-axis. Tolerant
plants will require more herbicide than non tolerant like plants in order to produce a given
herbicidal effect. Plants which are substantially "resistant" to the herbicide exhibit few, if
any, necrotic, lytic, chlorotic or other lesions when subjected to the herbicide at
15 concentrations and rates which are typically employed by the agrochemical community to
kill weeds in the field. Plants which are resistant to a herbicide are also tolerant of the
herbicide. The terms "resistant" and "tolerant" are to be construed as "tolerant and/or
resistant" within the context of the present application.

The herbicides of particular relevance to the present invention are those which are
20 capable *in vitro* of inhibiting 4-Hydroxy-phenylpyruvate dioxygenase (HPPD or 4HPPD)
enzymes. Such herbicides have been disclosed, such as the isoxazoles described especially in
the French Patent Applications 95 06800 and 95 13570 and especially isoxaflutole, a
selective maize herbicide, diketonitriles such as those described in European Applications 0
496 630, 0496 631, in particular 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-CF₃-
25 phenyl)propane-1,3-dione and 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-2,3Cl₂phenyl)propane-
1,3-dione, triketones described in European Applications 0 625 505 and 0 625 508, in
particular sulcotrione, mesotrione (BSI-proposed), pyrazolynate and pyrazoxyfen. Known
genes capable of providing for tolerance to these herbicides are those which encode HPPD
enzymes.

According to the present invention there is provided a method of making plants which are resistant or tolerant to herbicides which - *in vitro* - inhibit 4-hydroxyphenylpyruvate dioxygenase (4HPPD) comprising the steps of:

(i) transforming plant material with a polynucleotide comprising a region encoding a phytoene desaturase (PDS);

(ii) regenerating the thus transformed material into morphologically normal plants.

The region comprised by the polynucleotide may have the sequence depicted in SEQ ID No. 1, or may be a sequence which is complementary to one which when incubated at a temperature of between 55 and 60°C in 0.3 strength citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered saline containing 0.1% SDS still hybridises with the sequence depicted in SEQ ID No. 1.

It is preferred that the phytoene desaturase is of bacterial origin such as that depicted in SEQ ID No. 1 and being derived from *Erwinia uredovora*, and/or in particular is one which does not require plastoquinone 9 as a co-factor. The desaturase may, however be of plant origin, such as especially of monocotyledonous or dicotyledonous plants, especially of *Arabidopsis* or of *Umbelliferae*, such as, for example, the carrot (*Daucus carotta*). It can be native or possibly mutated while at the same time fundamentally retaining a property of herbicidal tolerance against HPPD inhibitors, such as herbicides of the isoxazoles family such as the Balance™ Herbicide or triketones. The herbicide resistant plants produced by the above method may be selected through their resistance to herbicides which *in vitro*, inhibit 4HPPD.

In may however, be further preferred that the polynucleotide encoding the phytoene desaturase further comprises a selectable marker gene to facilitate the selection of regenerated transformants. Suitable selectable marker genes include; resistance to antibiotics such as kanamycin, hygromycin and gentamycin; resistance to further herbicides such as glyphosate based herbicides; resistance to toxins such as eutypine.

Other forms of selection are also available such as hormone based selection systems such as the Multi Auto Transformation (MAT) system of Hiroyasu Ebinuma *et al.* 1997. PNAS Vol. 94 pp2117-2121; visual selection systems which use the known green fluorescence protein, β glucuronidase, mannose isomerase, xylose isomerase and 2-DOG.

The plant material may be, or may have been, further transformed with a polynucleotide comprising a region encoding a protein capable of providing the plant material with

resistance or tolerance to herbicides, insects, desiccation and/or fungal, bacterial or viral infections, or with a polynucleotide capable of encoding proteins which provide for improved quality traits such as increased yield, altered starch quality and/or increased nutrient content.

5 The protein encoding sequences within the polynucleotide are bounded by plant operable promoters and terminators. Such promoters and terminators, which are *per se* not germane to the invention, are well known to the skilled man and include, for example, the CaMV35S, FMV35S, NOS, OCS and E9 (derived from the small subunit of RUBISCO) promoters and terminators, or the promoter and terminator of a gene of α -tubulin (EP-A
10 652,286). Preferably, recourse is made to a promoter regulation sequence which favours the over-expression of the coding sequence, such as, for example, that comprising at least one histone promoter such as described in EP-A-507,698.

According to the invention, it is equally possible to use, in association with the promoter regulation sequence, other regulation sequences which are situated between the
15 promoter and the coding sequence, such as transcriptional or translational enhancers such as, for example, tobacco etch virus (TEV) translation activator described in International Patent application, PCT publication number WO87/07644 which is incorporated herein by reference, or of transit peptides, either single, or double, and in this case possibly separated by an intermediate sequence, that is to say comprising, in the transcription direction, a
20 sequence coding for a transit peptide of a plant gene coding for a plastid localization enzyme, a part of the sequence of the N-terminal mature part of a plant gene coding for a plastid localization enzyme, then a sequence coding for a second transit peptide of a plant gene coding for a plastid localization enzyme, formed by a part of the sequence of the N-terminal mature part of a plant gene coding for a plastid localization enzyme, such as described in EP-
25 A-508,909.

The plant material may have been, or may subsequently be - further transformed with a polynucleotide comprising a region encoding a protein capable of providing the plant with resistance or tolerance to herbicides, insects, desiccation and/or fungal, bacterial or viral infections, or with a polynucleotide capable of encoding proteins which provide for
30 improved quality traits such as increased yield, altered starch quality and/or increased nutrient content.

The protein capable of providing for herbicide resistance may be selected from the group consisting of glyphosate oxido-reductase (GOX), 5-enol-pyruvyl-3-phosphoshikimate synthetase (EPSPS), phosphinothricin acetyl transferase (PAT), hydroxyphenyl pyruvate dioxygenase (HPPD), glutathione S transferase (GST), cytochrome P450, Acetyl-CoA carboxylase (ACCase), Acetolactate synthase (ALS), protoporphyrinogen oxidase (PROTOX), dihydropteroate synthase, polyamine transport proteins, superoxide dismutase (SOD), bromoxynil nitrilase, the product of the *tfdA* gene obtainable from *Alcaligenes eutrophus*, and known mutagenised or otherwise modified variants of the said proteins.

As indicated above, the polynucleotide with which the plant material may be transformed may comprise 5' of the protein encoding regions which encode: (i) a peptide which is capable of targeting the translation products of the regions to plastids such as chloroplasts, mitochondria, other organelles or plant cell walls; and/or (ii) non-translated translational enhancing sequences.

The polynucleotide may be codon-optimised, or otherwise altered to enhance at least transcription once it is incorporated into plant material. Thus the polynucleotide used to transform the material may be modified in that mRNA instability encoding motifs and/or fortuitous splice regions may be removed, or plant preferred codons may be used so that expression of the thus modified polynucleotide in a plant yields substantially similar protein having a substantially similar activity/function to that obtained by expression of the unmodified polynucleotide in the organism in which the protein encoding regions of the unmodified polynucleotide are endogenous, with the *proviso* that if - in respect of the herbicide resistance conferring regions - the thus modified polynucleotide comprises plant preferred codons, the degree of identity between the protein encoding regions within the modified polynucleotide and like protein encoding regions endogenously contained within the said plant and encoding substantially the same protein is less than about 70%.

Transformation techniques are well known and include particle mediated biolistic transformation, *Agrobacterium*-mediated transformation, protoplast transformation (optionally in the presence of polyethylene glycols); sonication of plant tissues, cells or protoplasts in a medium comprising the polynucleotide or vector; micro-insertion of the polynucleotide or vector into totipotent plant material (optionally employing the known silicon carbide "whiskers" technique), electroporation and the like.

The invention still further provides morphologically normal fertile (or male sterile) whole plants regenerated from the material mentioned in the paragraph immediately preceding the last and the progeny of such plants, the seed of such plants and progeny, and parts of such plants and progeny. The transformed inventive plants include small grain cereals, oil seed
5 crops, fibre plants, fruit, vegetables, plantation crops and trees. Particularly preferred such plants include soybean, cotton, tobacco, sugarbeet, oilseed rape, canola, flax, sunflower, potato, tomato, alfalfa, lettuce, maize, wheat, sorghum, rye, bananas, barley, oat, turf grass, forage grass, sugar cane, pea, field bean, rice, pine, poplar, apple, grape, citrus and nut plants.

The transformed plants of the invention have tolerance or resistance to certain
10 herbicides such as the isoxazoles described especially in French Patent Applications 9506800 and 95 13570 and especially of 4-[4-CF₃-2-(methylsulphonyl)benzoyl]-5-cyclopropylisoxazole, and especially isoxaflutole, a selective maize herbicide, the diketetonitriles such as those described in EP-A-496,630 and EP-A-496,631, in particular 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-CF₃-phenyl)propane-1,3-dione and 2-cyano-3-
15 cyclopropyl-1-(2-SO₂CH₃-4-2,3-Cl₂-phenyl)propane-1,3-dione, and the triketones described in EP-A-625,505 and EP-A-625,508, in particular sulcotrione, mesotrione (BSI-proposed), pyrazolynate and pyrazoxyfen.

The invention further includes a morphologically normal fertile (or male sterile) whole plant resulting from the method of the invention, the progeny of such plants, the seed
20 of such plants and progeny, and parts of such plants and progeny.

The invention still further provides the use of a polynucleotide comprising a region encoding a phytoene desaturase in the production of plant material which is resistant or tolerant to herbicides which - *in vitro* - inhibit the enzyme 4-HPPD.

The invention still further provides a method of selectively controlling weeds in a
25 field, the field comprising weeds and crop plants, the method comprising application to the field of a herbicide which - *in vitro* - is capable of inhibiting the enzyme 4-HPPD, characterised in that the plants have been transformed with and express the coding regions of a polynucleotide comprising a sequence encoding a phytoene desaturase.

It is particularly preferred that the phytoene desaturase encoding sequence is that
30 which is depicted in SEQ ID No. 1, or is complementary to one which when incubated at a temperature of between 55 and 60°C in 0.3 strength citrate buffered saline containing 0.1%

SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered saline containing 0.1% SDS still hybridises with the sequence depicted in SEQ ID No. 1. The herbicide may be selected from the group consisting of mesotrione (BSI-proposed), pyrazolynate and pyrazoxyfen, Balance,TM sulcotrione etc. The field may be treated with a pesticide selected from the group consisting of a fungicide, insecticide and nematicide, either prior to or post application to the field of the herbicide.

The invention will now be described by way of the following non-limiting example, figure and the Sequence Listing in which:

SEQ ID No. 1 is the sequence of the phytoene desaturase (dehydrogenase) gene isolated from *Erwinia uredovora*. The person skilled in the art will recognise that any phytoene desaturase gene may be used in the production of plants having resistance/tolerance to the herbicides described above.

SEQ ID No. 2 is the protein encoded by SEQ ID No 1.

SEQ ID No.3 is the polynucleotide sequence encoding the pea rubisco small subunit transit peptide.

SEQ IN No. 4 is the amino acid sequence encoded by SEQ ID No. 3.

Figure 1 is the structure of plasmid pYPEIT4 carrying the *Erwinia uredovora* crtI gene with the transit peptide sequence (depicted as TP) of the pea rubisco small subunit.

EXAMPLE

Production of plants tolerant to herbicides capable of inhibiting the enzyme 4-HPPD *in vitro*.

The PDS gene (crtI) was cloned from *Erwinia uredovora*, a non-green phytopathogenic bacterial rot, and over-expressed in transgenic tobacco and tomato using a plasmid containing the CaMV 35S promoter and a chloroplast transit peptide (pYPIET4) (Misawa *et al.*, 1993). Homozygous tomato lines over-expressing the crtI gene were obtained as were tobacco plants containing the same construct.

Construction of plasmid pYPIET4 carrying the tp-crtI gene

Recombinant DNA techniques were performed using standard methods. A DNA sequence coding for the transit peptide (TP) in the precursor of the ribulose-1,5-bisphosphate carboxylase (Rubisco) small subunit of pea was isolated from plasmid pSNIF83 (Schreier *et al.*, 1985) as a 204 bp HindIII-SphI fragment, whose SphI site contains the tp processing site.

5 Plasmid pCRT-1 (Fraser *et al.* (1992) J.Biol.Chem 267 19891-19895) carrying the intact phytoene desaturase gene (*crtI*) of *Erwinia uredovora* was digested with BamHI and HindIII, and a 1.57 kb BamHI-HindIII fragment carrying the truncated *crtI* gene was isolated. The above 204 bp HindIII-SphI TP fragment was ligated with a 76bp synthesized fragment which carries the reading frame from the cohesive end for the SphI site containing the *crtI*

10 initiation codon to that of the BamHI site, and with the 1.57 kb BamHI-HindIII fragment. The desired 1.84 kb HindIII fragment carrying the tp-*crtI* chimeric gene was isolated, filled in with Klenow enzyme, and ligated into the SmaI-SacI site of a 10.9 kb fragment removing the β -glucuronidase gene from the binary vector bB/121 (purchased from Clontech laboratories). Thus, the desired plasmid pYPEIT4 was created, shown in Figure 1. The initiation codons

15 for the transit peptide and the intact *CrtI* are underlined. This HindIII fragment carrying the tp-*crtI* gene is surrounded by the CaMV 35S promoter and the NOS terminator of the binary vector pBI121 in order to lead to sufficient expression in the tissues of transgenic tobacco and tomato plants. As a control, plasmid pBICAR4 was constructed which carries an intact *crtI* gene without tp surrounded by the CaMV 35S promoter and the NOS terminator. The

20 plasmid pYPEIT4 was introduced into tobacco and tomato material by known techniques and the material then regenerated into intact plants, again by known techniques.

Tolerance of Tomato Plants transformed with *crtI* gene to Mesotrione and Isoxaflutole

Homozygous seed of tomato plants cv. Ailsa Craig, derived from 'wild type' (i.e. un-

25 transformed) and plants transformed with the *crtI* gene from *Erwinia uredovora*, (see above) were sown in a peat-based compost in 3 inch pots and transferred to the glasshouse. Plants were grown at 20/16 degrees day/night temperature under a 16 hour photoperiod for approximately 4 weeks prior to post-emergence treatment of four replicates with mesotrione or isoxaflutole (Balance TM Herbicide) at the 3 leaf stage. The chemicals were suspended in

30 water and applied, via a track sprayer at a spray volume of 200 litres per hectare, at rates ranging from 1 to 500 grammes active ingredient per hectare (g a.i./ha), as shown in Table 1.

The plants were left to grow for a further 25 days and then assessed visually for herbicidal damage compared to untreated 'control' plants. Typical phytotoxic symptoms observed were extreme chlorosis/bleaching and necrosis of leaves and new growth. The results from this test are given in Table 1 below where the '% Damage/Phytotoxicity' scores represent the mean of the visual assessment from each of the four treatment replicates.

Table 1

Chemical	Rate (g a.i./ha)	% Damage/Phytotoxicity (25 days after treatment)	
		Wild Type (Un-Transformed)	Transformed (crtI)
Mesotrione	1	25	0
	3	29	0
	11	50	9
	33	81	49
Isoxaflutole (Balance™)	1	11	2
	5	15	4
	15	19	6
	50	21	8
	150	34	19
	500	65	31

- As can be seen, plants transformed with the crtI gene which expresses the bacterial PDS from
- 5 *Erwinia uredovora*, demonstrate elevated tolerance to mesotrione and isoxaflutole compared to wild type, un-transformed tomatoes. For example, 11 g a.i./ha of mesotrione caused 50% phytotoxicity to wild type tomatoes but only 9% injury is observed in the transformed plants. Similarly, wild type plants are significantly more damaged by 500 g a.i./ha of isoxaflutole than those containing the crtI gene.
- 10 The skilled man will recognise that the invention is not limited to that described above. For example, plants other than tomato and tobacco may be transformed with a gene encoding a PDS enzyme, whether derived from a bacterial source or otherwise.

CLAIMS


1. A method of making plants which are resistant or tolerant to herbicides which - *in vitro* - inhibit 4-hydroxyphenylpyruvate dioxygenase (4HPPD) comprising the steps of:
 - (i) transforming plant material with a polynucleotide comprising a region encoding a phytoene desaturase (PDS);
 - (ii) regenerating the thus transformed material into morphologically normal plants and selecting from the population of regenerants those plants which are resistant or tolerant to herbicides which *in vitro* inhibit 4HPPD.
2. A method according to claim 1, wherein the region comprised by the polynucleotide is the sequence depicted in SEQ ID No. 1, or is a sequence which is complementary to one which when incubated at a temperature of between 55 and 60°C in 0.3 strength citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.3 strength citrate buffered saline containing 0.1% SDS still hybridises with the sequence depicted in SEQ ID No. 1.
3. A method according to claim 1, wherein the phytoene desaturase is of plant origin.
4. A method according to claim 1, wherein the phytoene desaturase is of bacterial origin.
5. A method according to claim 4 wherein the phytoene desaturase is isolatable from *Erwinia uredovora*.
6. A method according to any one of claims 1 to 5 wherein the polynucleotide further comprises a selectable marker gene.

7. A method according to claim 6 wherein the said selectable marker gene is selected from the group consisting of antibiotic resistance conferring, herbicide resistance conferring, toxin resistance conferring, nutritional markers, visual markers and marker genes used in hormone based selection systems.
- 5
8. A method according to any one of claims 1 to 7, wherein the plant material has been or is further transformed with a polynucleotide comprising a region encoding a protein capable of providing the plant material with resistance or tolerance to herbicides, insects, desiccation and/or fungal, bacterial or viral infections, or with a polynucleotide capable of encoding proteins which provide for improved quality traits such as increased yield, altered starch quality and/or increased nutrient content.
- 10
9. A method according to any one of claims 1 to 8, wherein the protein encoding sequences within the polynucleotide are bounded by plant operable promoters and terminators.
- 15
10. A method according to either of claims 8 or 9, wherein the protein capable of providing for herbicide resistance is selected from the group consisting of glyphosate oxido-reductase (GOX), 5-enol-pyruvyl-3-phosphoshikimate synthetase (EPSPS), phosphinothricin acetyl transferase (PAT), hydroxyphenyl pyruvate dioxygenase (HPPD), glutathione S transferase (GST), cytochrome P450, Acetyl-CoA carboxylase (ACCase), Acetolactate synthase (ALS), protoporphyrinogen oxidase (PROTOX), dihydropteroate synthase, polyamine transport proteins, superoxide dismutase (SOD), bromoxynil nitrilase, the product of the *tfdA* gene obtainable from *Alcaligenes eutrophus*, farnesyl pyrophosphate synthase and known mutagenised or otherwise modified variants of the said proteins.
- 20
- 25

11. A method according to any one of claims 1 to 10, wherein the protein encoding sequences of the polynucleotide comprise 5' regions which encode: (i) a peptide which is capable of targeting the translation products of the regions to plastids such as chloroplasts, mitochondria, other organelles or plant cell walls; and/or (ii) non-translated translational enhancing sequences.
12. A method according to any one of claims 1 to 11, in which the polynucleotide used to transform the material is modified in that mRNA instability encoding motifs and/or fortuitous splice regions are removed, or plant preferred codons are used so that expression of the thus modified polynucleotide in a plant yields substantially similar protein having a substantially similar activity/function to that obtained by expression of the unmodified polynucleotide in the organism in which the protein encoding regions of the unmodified polynucleotide are endogenous, with the *proviso* that if - in respect of the herbicide resistance conferring regions - the thus modified polynucleotide comprises plant preferred codons, the degree of identity between the protein encoding regions within the modified polynucleotide and like protein encoding regions endogenously contained within the said plant and encoding substantially the same protein is less than about 70%.
13. A method according to any one of claims 1 to 12, in which the 4-HPPD inhibiting herbicide is selected from the group consisting of isoxaflutole, diketonitriles such as 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-CF₃-phenyl)propane-1,3-dione and 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-2,3Cl₂-phenyl)propane-1,3-dione, triketones such as sulcotrione, and mesotrione (BSI-proposed), pyrazolynate and pyrazoxyfen.
14. A method according to any one of claims 1 to 13, wherein the herbicide is applied post-germination.
15. A morphologically normal fertile (or male sterile) whole plant resulting from the method of any one of claims 1 to 14, the progeny of such plants, the seed of such plants and progeny, and parts of such plants and progeny.

16. A plant according to claim 15 selected from the group consisting of banana, cotton, maize, tomato, vines.
- 5 17. Use of a polynucleotide comprising a region encoding a phytoene desaturase in the production of plant material which is resistant or tolerant to herbicides which - *in vitro* - inhibit the enzyme 4-HPPD.
18. A method of selectively controlling weeds in a field, the field comprising weeds and
10 crop plants, the method comprising application to the field of a herbicide which - *in vitro* - is capable of inhibiting the enzyme 4-HPPD, characterised in that the plants have been transformed with and express the coding regions of a polynucleotide comprising a sequence encoding a phytoene desaturase.
- 15 19. A method according to claim 18 wherein the polynucleotide is that mentioned in any one of claims 2 to 12.
20. A method according to either of claims 18 or 19, wherein the said herbicide is selected from the group consisting of, isoxaflutole, diketonitriles such as 2-cyano-3-
20 cyclopropyl-1-(2-SO₂CH₃-4-CF₃-phenyl)propane-1,3-dione and 2-cyano-3-cyclopropyl-1-(2-SO₂CH₃-4-2,3Cl₂phenyl)propane-1,3-dione, triketones such as sulcotrione, and mesotrione (BSI-proposed), pyrazolynate and pyrazoxyfen.
21. A method according to any one of claims 18 to 20, wherein the field is treated with a
25 pesticide selected from the group consisting of a fungicide, insecticide and nematocide, either prior to or post application to the field of the herbicide.

1/1

FIGURE 1

SEQUENCE LISTING

<110> ZENECA LIMITED

<120> METHOD OF PRODUCING PLANTS WHICH ARE TOLERANT OR
RESISTANT TO HERBICIDES

<130> PPD50336WO

<140>

<141>

<150> 9807818.1

<151> 1998-04-09

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 1493

<212> DNA

<213> Erwinia uredovora

<220>

<221> CDS

<222> (15)..(1493)

<400> 1

taaagagcgac	ctac	atg	aaa	cca	act	acg	gta	att	ggt	gca	ggc	ttc	ggt	50	
		Met	Lys	Pro	Thr	Thr	Val	Ile	Gly	Ala	Gly	Phe	Gly		
		1				5					10				
ggc	ctg	gca	ctg	gca	att	cgt	cta	caa	gct	gcg	ggg	atc	ccc	gtc	50
Gly	Leu	Ala	Leu	Ala	Ile	Arg	Leu	Gln	Ala	Ala	Gly	Ile	Pro	Val	50
	15					20					25				
ctg	ctt	gaa	caa	cgt	gat	aaa	ccc	ggc	ggt	cgg	gct	tat	gtc	tac	146
Leu	Leu	Glu	Gln	Arg	Asp	Lys	Pro	Gly	Gly	Arg	Ala	Tyr	Val	Tyr	146
	30					35				40					
gat	cag	ggg	ttt	acc	ttt	gat	gca	ggc	ccg	acg	ggt	atc	acc	gat	194
Asp	Gln	Gly	Phe	Thr		Phe	Asp	Ala	Gly	Pro	Thr	Val	Ile	Thr	194
	45				50					55				60	
agt	gcc	att	gaa	gaa	ctg	ttt	gca	ctg	gca	gga	aaa	cag	tta	aaa	242
Ser	Ala	Ile	Glu	Glu	Leu	Phe	Ala	Leu	Ala	Gly	Lys	Gln	Leu	Lys	242
			65					70				75			
tat	gtc	gaa	ctg	ctg	ccg	ggt	acg	ccg	ttt	tac	cgc	ctg	tgt	tgg	290
Tyr	Val	Glu	Leu	Leu	Pro	Val	Thr	Pro	Phe	Tyr	Arg	Leu	Cys	Trp	290
		80					85					90			
tca	ggg	aag	gtc	ttt	aat	tac	gat	aac	gat	caa	acc	cgg	ctc	gaa	338
Ser	Gly	Lys	Val	Phe	Asn	Tyr	Asp	Asn	Asp	Gln	Thr	Arg	Leu	Glu	338
	95						100					105			
cag	att	cag	cag	ttt	aat	ccc	cgc	gat	gtc	gaa	ggt	tat	cgt	cag	386
Gln	Ile	Gln	Gln	Phe	Asn	Pro	Arg	Asp	Val	Glu	Gly	Tyr	Arg	Gln	386
	110					115				120					
ctg	gac	tat	tca	cgc	gcg	gtg	ttt	aaa	gaa	ggc	tat	cta	aag	ctc	434
Leu	Asp	Tyr	Ser	Arg	Ala	Val	Phe	Lys	Glu	Gly	Tyr	Leu	Lys	Leu	434
	125				130					135			140		

-2-

act gtc cct ttt tta tcg ttc aga gac atg ctt cgc gcc gca cct caa	482
Thr Val Pro Phe Leu Ser Phe Arg Asp Met Leu Arg Ala Ala Pro Gln	
145 150 155	
ctg gcg aaa ctg cag gca tgg aga agc gtt tac agt aag gtt gcc agt	530
Leu Ala Lys Leu Gln Ala Trp Arg Ser Val Tyr Ser Lys Val Ala Ser	
160 165 170	
tac atc gaa gat gaa cat ctg cgc cag gcg ttt tct ttc cac tcg ctg	578
Tyr Ile Glu Asp Glu His Leu Arg Gln Ala Phe Ser Phe His Ser Leu	
175 180 185	
ttg gtg ggc ggc aat ccc ttc gcc acc tca tcc att tat acg ttg ata	626
Leu Val Gly Gly Asn Pro Phe Ala Thr Ser Ser Ile Tyr Thr Leu Ile	
190 195 200	
cac gcg ctg gag cgt gag tgg ggc gtc tgg ttt cgc cgt ggc gcc acc	674
His Ala Leu Glu Arg Glu Trp Gly Val Trp Phe Pro Arg Gly Gly Thr	
205 210 215 220	
ggc gca tta gtt cag ggg atg ata aag ctg ttt cag gat ctg ggt ggc	722
Gly Ala Leu Val Gln Gly Met Ile Lys Leu Phe Gln Asp Leu Gly Gly	
225 230 235	
gaa gtc gtg tta aac gcc aga gtc agc cat atg gaa acg aca gga aac	770
Glu Val Val Leu Asn Ala Arg Val Ser His Met Glu Thr Gly Asn	
240 245 250	
aag att gaa gcc gtg cat tta gag gac ggt cgc agg ttc ctg acg caa	818
Lys Ile Glu Ala Val His Leu Glu Asp Gly Arg Arg Phe Leu Thr Gln	
255 260 265	
gcc gtc gcg tca aat gca gat gtg gtt cat acc tat cgc gac ctg tta	866
Ala Val Ala Ser Asn Ala Asp Val Val His Thr Tyr Arg Asp Leu Leu	
270 275 280	
agc cag cac cct gcc gcg gtt aag cag tcc aac aaa ctg cag act aag	914
Ser Gln His Pro Ala Ala Val Lys Gln Ser Asn Lys Leu Gln Thr Lys	
285 290 295 300	
cgc atg agt aac tct ctg ttt gtg ctc tat ttt ggt ttg aat cac cat	962
Arg Met Ser Asn Ser Leu Phe Val Leu Tyr Phe Gly Leu Asn His His	
305 310 315	
cat gat cag ctc gcg cat cac acg gtt tgt ttc gcc ccg cgt tac cgc	1010
His Asp Gln Leu Ala His His Thr Val Cys Phe Gly Pro Arg Tyr Arg	
320 325 330	
gag ctg att gac gaa att ttt aat cat gat ggc ctc gca gag gac ttc	1058
Glu Leu Ile Asp Glu Ile Phe Asn His Asp Gly Leu Ala Glu Asp Phe	
335 340 345	
tca ctt tat ctg cac gcg ccc tgt gtc acg gat tcg tca ctg gcg cct	1106
Ser Leu Tyr Leu His Ala Pro Cys Val Thr Asp Ser Ser Leu Ala Pro	
350 355 360	
gaa ggt tgc ggc agt tac tat gtg ttg gcg ccg gtg ccg cat tta gcc	1154
Glu Gly Cys Gly Ser Tyr Tyr Val Leu Ala Pro Val Pro His Leu Gly	
365 370 375 380	
acc gcg aac ctc gac tgg acg gtt gag ggg cca aaa cta cgc gac cgt	1202
Thr Ala Asn Leu Asp Trp Val Glu Gly Pro Lys Leu Arg Asp Arg	
385 390 395	
att ttt gcg tac ctt gag cag cat tac atg cct gcc tta cgg agt cag	1250
Ile Phe Ala Tyr Leu Glu Gln His Tyr Met Pro Gly Leu Arg Ser Gln	

-3-

400	405	410	
ctg gtc acg cac cgg atg ttt acg ccg ttt gat ttt cgc gac cag ctt			1298
Leu Val Thr His Arg Met Phe Thr Pro Phe Asp Phe Arg Asp Gln Leu			
415	420	425	
aat gcc tat cat ggc tca gcc ttt tct gtg gag ccc gtt ctt acc cag			1346
Asn Ala Tyr His Gly Ser Ala Phe Ser Val Glu Pro Val Leu Thr Gln			
430	435	440	
agc gcc tgg ttt cgg ccg cat aac cgc gat aaa acc att act aat ctc			1394
Ser Ala Trp Phe Arg Pro His Asn Arg Asp Lys Thr Ile Thr Asn Leu			
445	450	455	460
tac ctg gtc ggc gca ggc acg cat ccc gcc gca gcc att cct gcc gtc			1442
Tyr Leu Val Gly Ala Gly Thr His Pro Gly Ala Gly Ile Pro Gly Val			
465	470	475	
atc gcc tcg gca aaa gcg aca gca ggt ttg atg ctg gag gat ctg att			1490
Ile Gly Ser Ala Lys Ala Thr Ala Gly Leu Met Leu Glu Asp Leu Ile			
480	485	490	
tga			1493
<210> 2			
<211> 492			
<212> PRT			
<213> Erwinia uredovora			
<400> 2			
Met Lys Pro Thr Thr Val Ile Gly Ala Gly Phe Gly Gly Leu Ala Leu			
1	5	10	15
Ala Ile Arg Leu Gln Ala Ala Gly Ile Pro Val Leu Leu Leu Glu Gln			
20	25	30	
Arg Asp Lys Pro Gly Gly Arg Ala Tyr Val Tyr Glu Asp Gln Gly Phe			
35	40	45	
Thr Phe Asp Ala Gly Pro Thr Val Ile Thr Asp Pro Ser Ala Ile Glu			
50	55	60	
Glu Leu Phe Ala Leu Ala Gly Lys Gln Leu Lys Glu Tyr Val Glu Leu			
65	70	75	80
Leu Pro Val Thr Pro Phe Tyr Arg Leu Cys Trp Glu Ser Gly Lys Val			
85	90	95	
Phe Asn Tyr Asp Asn Asp Gln Thr Arg Leu Glu Ala Gln Ile Gln Gln			
100	105	110	
Phe Asn Pro Arg Asp Val Glu Gly Tyr Arg Gln Phe Leu Asp Tyr Ser			
115	120	125	
Arg Ala Val Phe Lys Glu Gly Tyr Leu Lys Leu Gly Thr Val Pro Phe			
130	135	140	
Leu Ser Phe Arg Asp Met Leu Arg Ala Ala Pro Gln Leu Ala Lys Leu			
145	150	155	160
Gln Ala Trp Arg Ser Val Tyr Ser Lys Val Ala Ser Tyr Ile Glu Asp			
165	170	175	
Glu His Leu Arg Gln Ala Phe Ser Phe His Ser Leu Leu Val Gly Gly			

-4-

180	185	190
Asn Pro Phe Ala Thr Ser Ser Ile Tyr Thr Leu Ile His Ala Leu Glu		
195	200	205
Arg Glu Trp Gly Val Trp Phe Pro Arg Gly Gly Thr Gly Ala Leu Val		
210	215	220
Gln Gly Met Ile Lys Leu Phe Gln Asp Leu Gly Gly Glu Val Val Leu		
225	230	235
Asn Ala Arg Val Ser His Met Glu Thr Thr Gly Asn Lys Ile Glu Ala		
	245	250
Val His Leu Glu Asp Gly Arg Arg Phe Leu Thr Gln Ala Val Ala Ser		
	260	265
Asn Ala Asp Val Val His Thr Tyr Arg Asp Leu Leu Ser Gln His Pro		
	275	280
Ala Ala Val Lys Gln Ser Asn Lys Leu Gln Thr Lys Arg Met Ser Asn		
	290	295
Ser Leu Phe Val Leu Tyr Phe Gly Leu Asn His His His Asp Gln Leu		
305	310	315
Ala His His Thr Val Cys Phe Gly Pro Arg Tyr Arg Glu Leu Ile Asp		
	325	330
Glu Ile Phe Asn His Asp Gly Leu Ala Glu Asp Phe Ser Leu Tyr Leu		
	340	345
His Ala Pro Cys Val Thr Asp Ser Ser Leu Ala Pro Glu Gly Cys Gly		
	355	360
Ser Tyr Tyr Val Leu Ala Pro Val Pro His Leu Gly Thr Ala Asn Leu		
	370	375
Asp Trp Thr Val Glu Gly Pro Lys Leu Arg Asp Arg Ile Phe Ala Tyr		
385	390	395
Leu Glu Gln His Tyr Met Pro Gly Leu Arg Ser Gln Leu Val Thr His		
	405	410
Arg Met Phe Thr Pro Phe Asp Phe Arg Asp Gln Leu Asn Ala Tyr His		
	420	425
Gly Ser Ala Phe Ser Val Glu Pro Val Leu Thr Gln Ser Ala Trp Phe		
	435	440
Arg Pro His Asn Arg Asp Lys Thr Ile Thr Asn Leu Tyr Leu Val Gly		
	450	455
Ala Gly Thr His Pro Gly Ala Gly Ile Pro Gly Val Ile Gly Ser Ala		
465	470	475
Lys Ala Thr Ala Gly Leu Met Leu Glu Asp Leu Ile		
	485	490

<210> 3

<211> 204

<212> DNA

<213> Pisum sativum

<220>

-5-

<221> CDS

<222> (1)..(204)

<400> 3

atg gct tct atg ata tcc tct tcg gct gtg aca aca gtc agc cgt gcc	48
Met Ala Ser Met Ile Ser Ser Ser Ala Val Thr Val Ser Arg Ala	
1 5 10 15	

tct agg ggg caa tcc gcc gca gtg gct cca ttc ggc ggc ctc aaa tcc	96
Ser Arg Gly Gln Ser Ala Ala Val Ala Pro Phe Gly Gly Leu Lys Ser	
20 25 30	

atg act gga ttc cca gtg aag aag gtc aac act gac att act tcc att	144
Met Thr Gly Phe Pro Val Lys Lys Val Asn Thr Asp Ile Thr Ser Ile	
35 40 45	

aca agc aat ggt gga aga gta aag tgc atg aaa cca act acg gta att	192
Thr Ser Asn Gly Gly Arg Val Lys Cys Met Lys Pro Thr Thr Val Ile	
50 55 60	

ggt gca ggc ttc	204
Gly Ala Gly Phe	
65	

<210> 4

<211> 68

<212> PRT

<213> Pisum sativum

<400> 4

Met Ala Ser Met Ile Ser Ser Ser Ala Val Thr Thr Val Ser Arg Ala	
1 5 10 15	

Ser Arg Gly Gln Ser Ala Ala Val Ala Pro Phe Gly Gly Leu Lys Ser	
20 25 30	

Met Thr Gly Phe Pro Val Lys Lys Val Asn Thr Asp Ile Thr Ser Ile	
35 40 45	

Thr Ser Asn Gly Gly Arg Val Lys Cys Met Lys Pro Thr Thr Val Ile	
50 55 60	

Gly Ala Gly Phe	
65	

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 A01H5/00 A01H5/10 //C12N15/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MISAWA N ET AL: "EXPRESSION OF AN ERWINA PHYTOENE DESATURASE GENE NOT ONLY CONFERS MULTIPLE RESISTANCE TO HERBICIDES INTERFERING WITH CAROTENOID BIOSYNTHESIS BUT ALSO ALTERS XANTHOPHYLL METABOLISM IN TRANSGENIC PLANTS" PLANT JOURNAL, vol. 6, no. 4, 1 January 1994 (1994-01-01), pages 481-489, XP002017203 ISSN: 0960-7412 the whole document	15, 16
A	page 486, column 1, paragraph 2; table 1	
A	WO 97 27285 A (UNIV ARIZONA) 31 July 1997 (1997-07-31) page 6, line 7-24 --- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X*" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y*" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

11 August 1999

Date of mailing of the international search report

30/08/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bilang, J

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BARTA I C ET AL: "BENZOYL-CYCLOHEXANEDIONE HERBICIDES ARE STRONG INHIBITORS OF PURIFIED P-HYDROXYPHENYLPYRUVIC ACID DIOXYGENASE OF MAIZE" PESTICIDE SCIENCE, vol. 45, no. 3, 1 November 1995 (1995-11-01), page 286/287 XP000547268 ISSN: 0031-613X the whole document ---</p>	
A	<p>SCHULZ A ET AL: "SC-0051, A 2-BENZOYL-CYCLOHEXANE-1,3-DIONE BLEACHING HERBICIDE, IS A POTENT INHIBITOR OF THE ENZYME P-HYDROXYPHENYLPYRUVATE DIOXYGENASE" FEBS LETTERS, vol. 318, no. 2, 1 March 1993 (1993-03-01), pages 162-166, XP002028049 ISSN: 0014-5793 abstract ---</p>	
A	<p>BABCZINSKI P ET AL: "SUBSTITUTED TETRAHYDROPYRIMIDINONES. A NEW CLASS OF HERBICIDAL COMPOUNDS INDUCING CHLOROSIS BY INHIBITION OF PHYTOENE DESATURATION" PESTICIDE SCIENCE, vol. 30, no. 3, 1 January 1990 (1990-01-01), pages 339-342, XP000202878 ISSN: 0031-613X the whole document ---</p>	
A	<p>MISAWA ET AL.: "E. uredovora carotenoid biosynthesis genes" EMBL DATABASE ACCESSION NUMBER D90087, 6 January 1991 (1991-01-06), XP002111545 the whole document -----</p>	

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 99/01059

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9727285 A	31-07-1997	AU 1845397 A EP 0877793 A	20-08-1997 18-11-1998
